

Supersymmetric Calogero models by gauging

Sergey Fedoruk¹, Evgeny Ivanov¹, Olaf Lechtenfeld²

¹*Bogoliubov Laboratory of Theoretical Physics, JINR, 141980 Dubna, Moscow Region, Russia*

²*Institut für Theoretische Physik, Leibniz Universität Hannover, Appelstraße 2, D-30167 Hannover, Germany*

New superconformal extensions of $d=1$ Calogero-type systems are obtained by gauging the $U(n)$ isometry of matrix superfield models. We consider the cases of $\mathcal{N}=1$, $\mathcal{N}=2$ and $\mathcal{N}=4$ one-dimensional supersymmetries. The bosonic core of the $\mathcal{N}=1$ and $\mathcal{N}=2$ models is the standard conformal A_{n-1} Calogero system, whereas the $\mathcal{N}=4$ model is an extension of the $U(2)$ -spin Calogero system.

PACS numbers: 11.30.Pb; 12.60.Jv; 02.30.Ik; 02.10.Yn

1. Introduction. Superconformal extensions of Calogero model [1] provide nice examples of integrable supersymmetric quantum-mechanical systems and as such are of vast interest from various points of view (see [2, 3] for the survey of physical applications of Calogero model). In particular, by a conjecture of Gibbons and Townsend [4], $\mathcal{N}=4$ superconformal models might be closely related to M-theory. While the $\mathcal{N}=2$ super Calogero models for any number of interacting particles were constructed in full generality rather long ago [5] (see also [6, 7]), until now there is no such an exhaustive understanding of the generic $\mathcal{N}=4$ models despite the existence of extensive literature on this subject (see e.g. [6, 7, 8, 9, 10, 11]). It seems important to develop some universal approach to superconformal Calogero-type models including the $\mathcal{N}=4$ ones.

The purpose of this letter is to present a candidate approach of this type suitable for an arbitrary number of interacting particles. It is based on the superfield gauging of some non-abelian isometries of the $d=1$ field theories. This gauging procedure was worked out in [12] to understand off-shell dualities between $d=1$ supermultiplets with different sets of physical bosonic components.

Our starting point is the nice interpretation of the bosonic n -particle Calogero model as a $U(n)$, $d=1$ gauge theory [13] (see also [14] and [15, 16]). In the formulation of [13] the model is described by the hermitian $n \times n$ -matrix field $X_a^b(t)$, $(\overline{X}_a^b) = X_b^a$, the complex n -plet $Z_a(t)$, $\bar{Z}^a = (\bar{Z}_a)$, $a, b = 1, \dots, n$, and n^2 non-propagating “gauge fields” $A_a^b(t)$, $(\bar{A}_a^b) = A_b^a$. The action reads

$$S_0 = \int dt \left[\text{Tr}(\nabla X \nabla X) + \frac{i}{2}(\bar{Z} \nabla Z - \nabla \bar{Z} Z) + c \text{Tr} A \right], \quad (1)$$

where the covariant derivatives are defined as

$$\nabla X = \dot{X} + i[A, X], \quad \nabla Z = \dot{Z} + iAZ.$$

The real constant c of the Calogero interaction comes out from a Fayet-Iliopoulos (FI) term in (1).

The action (1) is invariant with respect to the local $U(n)$ transformations, $g(\tau) \in U(n)$,

$$X \rightarrow gXg^\dagger, \quad Z \rightarrow gZ, \quad A \rightarrow gAg^\dagger + i\dot{g}g^\dagger, \quad (2)$$

and we can fully fix the $U(n)$ gauge freedom by choosing

$$X_a^b = x_a \delta_a^b, \quad \bar{Z}^a = Z_a. \quad (3)$$

Inserting these gauge conditions and the *algebraic* equations of motion $(Z_a)^2 = c$ (which implies $c > 0$) and $A_a^b = \frac{Z_a Z_b}{2(x_a - x_b)^2}$, $a \neq b$, back into the action (1), we arrive at the standard Calogero action

$$S_C = \int dt \left[\sum_a \dot{x}_a \dot{x}_a - \sum_{a \neq b} \frac{c^2}{4(x_a - x_b)^2} \right] \quad (4)$$

as a fixed gauge of (1). Note the important role of the auxiliary $U(n)$ multiplet Z with the $d=1$ Wess-Zumino (WZ) action in (1) for recovering the Calogero action.

The original action (1) is invariant under the $d=1$ conformal $SO(1,2)$ transformations: $\delta t = a$, $\delta X_a^b = \frac{1}{2} \dot{a} X_a^b$, $\delta Z_a = 0$, $\delta A_a^b = -\dot{a} A_a^b$, where $a(t)$ obeys the constraint $\ddot{a} = 0$. This property implies the well-known conformal invariance of the eventual Calogero model.

Our approach is a minimal superfield generalization of this bosonic $U(n)$ gauging. Requiring the supersymmetric gauge models to possess $\mathcal{N}$ -extended superconformal symmetry essentially constrains the structure of the corresponding actions and allows one to reveal, in their bosonic sector, either the standard Calogero model (4) (in the cases of $\mathcal{N}=1$ and $\mathcal{N}=2$) or the $U(2)$ -spin Calogero model [3, 17, 18] modified by a conformal potential for the center-of-mass coordinate (in the $\mathcal{N}=4$ case). In this short note we outline the basic features of our

construction, leaving details, quantization and comparison with the previously known superextended Calogero models for a longer paper.

2. $\mathcal{N}=1$ supersymmetric extension. We use the Grassmann-even hermitian $\mathcal{N}=1$ matrix superfield $\mathcal{X}_a^b(t, \theta)$, $(\mathcal{X})^\dagger = \mathcal{X}$, belonging to the adjoint representation of $U(n)$, as well as the Grassmann-even complex $\mathcal{N}=1$ superfield $\mathcal{Z}_a(t, \theta)$, $\bar{\mathcal{Z}}^a(t, \theta) = (\mathcal{Z}_a)^\dagger$, in the fundamental of $U(n)$. The spinor and time derivatives,

$$D = \partial_\theta + i\theta\partial_t, \quad \{D, D\} = 2i\partial_t,$$

are gauge-covariantized by the anti-hermitian Grassmann-odd connections $\mathcal{A}_a^b(t, \theta)$, $(\mathcal{A})^\dagger = -\mathcal{A}$:

$$D\mathcal{X} = D\mathcal{X} + i[\mathcal{A}, \mathcal{X}], \quad \nabla_t \mathcal{X} = -iD\mathcal{D}\mathcal{X}, \quad D\mathcal{Z} = D\mathcal{Z} + i\mathcal{A}\mathcal{Z}.$$

The minimal gauge invariant action has the following form ($\mu_1 = dt d\theta$):

$$S_1 = -i \int \mu_1 \left[\text{Tr} (\nabla_t \mathcal{X} D\mathcal{X} + c\mathcal{A}) + \frac{i}{2} (\bar{\mathcal{Z}} D\mathcal{Z} - D\bar{\mathcal{Z}} \mathcal{Z}) \right]. \quad (5)$$

It is invariant under the local $U(n)$ transformations:

$$\mathcal{X}' = e^{i\tau} \mathcal{X} e^{-i\tau}, \quad \mathcal{Z}' = e^{i\tau} \mathcal{Z}, \quad \mathcal{A}' = e^{i\tau} \mathcal{A} e^{-i\tau} - i e^{i\tau} D e^{-i\tau},$$

where $\tau_a^b(t, \theta) \in u(n)$ is the hermitian matrix parameter. It is also invariant under the $\mathcal{N}=1$ superconformal group. Conformal supersymmetry $\delta' t = -i\eta\theta t$, $\delta'\theta = \eta t$ has the following realization on the involved superfields:

$$\delta' \mathcal{X} = -i\eta\theta \mathcal{X}, \quad \delta' \mathcal{A} = i\eta\theta \mathcal{A}, \quad \delta' \mathcal{Z} = 0.$$

Its closure with the Poincaré $\mathcal{N}=1$, $d=1$ supersymmetry yields the full $\mathcal{N}=1$ superconformal symmetry.

Due to the $U(n)$ gauge invariance we can choose the WZ gauge for the spinor connection:

$$\mathcal{A} = i\theta \mathcal{A}(t). \quad (6)$$

Substituting this into the action (5), integrating there over θ and eliminating the auxiliary fields by their equations of motion, we obtain

$$S_1 = S_0 + S_1^\Psi, \quad S_1^\Psi = -i \text{Tr} \int dt \Psi \nabla \Psi \quad (7)$$

where $\Psi = -iD\mathcal{X}$ is the matrix Grassmann-odd field and $\nabla \Psi = \dot{\Psi} + i[\mathcal{A}, \Psi]$. The bosonic limit of (7) (and hence of (5)) is just the Calogero action (1). Its gauge $U(n)$ symmetry is the residual symmetry of the WZ gauge (6).

An alternative supersymmetric gauge choice is

$$\mathcal{X}_a^b = 0, \quad a \neq b \quad \Leftrightarrow \quad \mathcal{X}_a^b = \mathcal{X}_a \delta_a^b; \quad \mathcal{Z}_a = \bar{\mathcal{Z}}^a. \quad (8)$$

In this gauge the model is described by n^2 real $\mathcal{N}=1$ superfields $\mathcal{A}_a^b$, $a \neq b$, and $\mathcal{X}_a$ (the superfields $\mathcal{Z}_a$ and those on the diagonal of $\mathcal{A}_a^b$ are auxiliary). In the two-particle case ($n=2$) the resulting $\mathcal{N}=1$ system possesses

an additional hidden $\mathcal{N}=1$ supersymmetry, so that the $n=2$ model is in fact $\mathcal{N}=2$ superconformal mechanics plus an $\mathcal{N}=2$ free multiplet corresponding to the center-of-mass motion. Starting with $n=3$, one gets new $\mathcal{N}=1$ superextensions of the n -particle Calogero models which cannot be recovered by any truncations of the standard $\mathcal{N}=2$ superextensions [5].

3. $\mathcal{N}=2$ supersymmetric extension. The relevant superfield content consists of the hermitian matrix superfield $\mathcal{X}_a^b(t, \theta, \bar{\theta})$, $(\mathcal{X})^\dagger = \mathcal{X}$ with the off-shell field content $(\mathbf{1}, \mathbf{2}, \mathbf{1})$, and bosonic chiral $U(n)$ -fundamental superfield $\mathcal{Z}_a(t_L, \theta)$, $\bar{\mathcal{Z}}^a(t_R, \bar{\theta}) = (\mathcal{Z}_a)^\dagger$, $t_{L,R} = t \pm i\theta\bar{\theta}$,

$$\bar{D}\mathcal{Z}_a = 0, \quad D\bar{\mathcal{Z}}^a = 0, \quad (9)$$

with the field contents $(\mathbf{2}, \mathbf{2}, \mathbf{0})$. Here

$$D = \partial_\theta + i\bar{\theta}\partial_t, \quad \bar{D} = -\partial_{\bar{\theta}} - i\theta\partial_t, \quad \{D, \bar{D}\} = -2i\partial_t.$$

The gauge prepotential is an $n \times n$ hermitian matrix $V_a^b(t, \theta, \bar{\theta})$, $(V)^\dagger = V$. The action reads ($\mu_2 = dt d^2\theta$)

$$S_2 = \int \mu_2 \left[\text{Tr} (\bar{D}\mathcal{X} e^{2V} D\mathcal{X} e^{2V}) + \frac{1}{2} \bar{\mathcal{Z}} e^{2V} \mathcal{Z} - c \text{Tr} V \right], \quad (10)$$

the gauge-covariant derivatives being defined as

$$D\mathcal{X} = D\mathcal{X} + e^{-2V} (De^{2V}) \mathcal{X}, \quad \bar{D}\mathcal{X} = \bar{D}\mathcal{X} - \mathcal{X} e^{2V} (\bar{D}e^{-2V}).$$

The action (10) is invariant under the local $U(n)$ transformations:

$$\mathcal{X}' = e^{i\lambda} \mathcal{X} e^{-i\bar{\lambda}}, \quad \mathcal{Z}' = e^{i\lambda} \mathcal{Z}, \quad e^{2V'} = e^{i\bar{\lambda}} e^{2V} e^{-i\lambda} \quad (11)$$

where n^2 complex gauge parameters, $\lambda = (\lambda_a^b)$, are (anti)chiral superfields: $\lambda(t_L, \theta) \in u(n)$, $\bar{\lambda}(t_R, \bar{\theta}) = (\lambda)^\dagger \in u(n)$. The action is also invariant under the superconformal group $SU(1, 1|1)$. The conformal supersymmetry acts on the coordinates as

$$\delta' t = -i(\eta\bar{\theta} + \bar{\eta}\theta)t, \quad \delta'\theta = \eta(t + i\theta\bar{\theta})$$

and on the superfields as

$$\delta' \mathcal{X} = -i(\eta\bar{\theta} + \bar{\eta}\theta) \mathcal{X}, \quad \delta' V = 0, \quad \delta' \mathcal{Z} = 0.$$

The chirality conditions (9) are preserved by these transformations.

In the WZ gauge

$$V(t, \theta, \bar{\theta}) = -\theta\bar{\theta} \mathcal{A}(t) \quad (12)$$

the action (10) takes the form

$$S_2 = S_0 + S_2^\Psi, \quad S_2^\Psi = -i \text{Tr} \int dt (\bar{\Psi} \nabla \Psi - \nabla \bar{\Psi} \Psi) \quad (13)$$

where $\Psi = D\mathcal{X}$ is a Grassmann-odd field and

$$\nabla \Psi = \dot{\Psi} + i[\mathcal{A}, \Psi], \quad \nabla \bar{\Psi} = \dot{\bar{\Psi}} + i[\mathcal{A}, \bar{\Psi}]. \quad (14)$$

We see that the bosonic core of the action (13) exactly coincides with the Calogero action (1).

The action (13) is invariant with respect to the residual local bosonic $U(n)$ transformations, defined by (2) and $\Psi \rightarrow g\Psi g^\dagger$, therefore we can choose the gauge (3). As a result we obtain an $\mathcal{N}=2$ superextension of the n -particle Calogero model. In the two-particle case ($n=2$) we found that the $\mathcal{N}=2$ supersymmetric gauged system actually describes $\mathcal{N}=4$ superconformal mechanics plus one $\mathcal{N}=4$ free multiplet corresponding to the center-of-mass motion, so that there is a hidden extra $\mathcal{N}=2$ symmetry in this case. For $n > 2$ we obtain some new $\mathcal{N}=2$ extensions of the n -particle Calogero models with n bosonic variables and $n \times n$ fermionic ones, as opposed to the standard $\mathcal{N}=2$ super-Calogero with n complex fermions [5].

The presence of the matrix Grassmann-odd field Ψ in the action (13) (and also in (7)) is imperative for $d=1$ supersymmetry and superconformal symmetry. Similar structures with the bosonic analogs of Ψ appeared e.g. in [15, 16] in connection with the quantum Hall effect.

4. $\mathcal{N}=4$ supersymmetric extension. This case surprisingly yields $U(2)$ -spin Calogero system [17, 18] in the bosonic sector. The most natural formulation of $\mathcal{N}=4, d=1$ models is achieved in the harmonic superspace [19, 20] parametrized by the coordinates $(t, \theta_i, \bar{\theta}^k, u_i^\pm)$, $i, k = 1, 2$, where commuting mutually conjugate $SU(2)$ -doublets $u_i^\pm$ are harmonic coordinates, $u^{+i}u_i^- = 1$. The harmonic analytic subspace is parametrized by the coordinates $(\zeta, u) = (t_A, \theta^+, \bar{\theta}^+, u_i^\pm)$, $t_A = t - i(\theta^+\bar{\theta}^- + \theta^-\bar{\theta}^+)$, $\theta^\pm = \theta^i u_i^\pm$, $\bar{\theta}^\pm = \bar{\theta}^i u_i^\pm$. The integration measures are defined as $\mu_H = dudt d^4\theta$ and $\mu_A^{(-2)} = dud\zeta^{(-2)}$.

The $\mathcal{N}=4$ supersymmetric model with $U(n)$ gauge symmetry is described by the action

$$S_4 = S_X + S_{FI} + S_{WZ}. \quad (15)$$

The first term in (15)

$$S_X = -\frac{1}{2} \int \mu_H \text{Tr} (\mathcal{X}^2) \quad (16)$$

is the gauged action of the **(1,4,3)** multiplets. The latter are described by hermitian matrix superfields $\mathcal{X} = (\mathcal{X}_a^b)$ subjected to the gauge-covariant constraints

$$\mathcal{D}^{++} \mathcal{X} = 0, \quad (17)$$

$$\mathcal{D}^+ \mathcal{D}^- \mathcal{X} = 0, \quad (\mathcal{D}^+ \bar{\mathcal{D}}^- + \bar{\mathcal{D}}^+ \mathcal{D}^-) \mathcal{X} = 0. \quad (18)$$

The constraint (17) involves the covariant harmonic derivative $\mathcal{D}^{++} = D^{++} + iV^{++}$, where the gauge matrix connection $V^{++}(\zeta, u)$ is an analytic superfield.¹ The

gauge connections entering the spinor covariant derivatives in (18) are properly expressed through $V^{++}(\zeta, u)$ [12]. The parameters of the $U(n)$ gauge group are analytic, so $\mathcal{D}^+ = D^+$, $\bar{\mathcal{D}}^+ = \bar{D}^+$. Note that $\mathcal{X}$ is in the adjoint of $U(n)$, so $\mathcal{D}^{++} \mathcal{X} = D^{++} \mathcal{X} + i[V^{++}, \mathcal{X}]$, etc.

The second term in (15) is the FI term

$$S_{FI} = \frac{i}{2} c \int \mu_A^{(-2)} \text{Tr} V^{++}. \quad (19)$$

The third term in (15),

$$S_{WZ} = \frac{1}{2} \int \mu_A^{(-2)} \mathcal{V}_0 \tilde{\mathcal{Z}}^+ \mathcal{Z}^+, \quad (20)$$

is a WZ action describing n commuting analytic superfields $\mathcal{Z}_a^+$ (analogs of the superfields $\mathcal{Z}_a$ of the $\mathcal{N}=1$ and $\mathcal{N}=2$ cases). They represent off-shell $\mathcal{N}=4$ multiplets **(4,4,0)** and are defined by the constraints

$$\mathcal{D}^{++} \mathcal{Z}^+ = 0, \quad D^+ \mathcal{Z}^+ = 0, \quad \bar{D}^+ \mathcal{Z}^+ = 0. \quad (21)$$

At last, the superfield $\mathcal{V}_0(\zeta, u)$ is a real analytic gauge superfield, $D^+ \mathcal{V}_0 = 0$, $\bar{D}^+ \mathcal{V}_0 = 0$, which is defined by the integral transform [12]

$$\mathcal{X}_0(t, \theta_i, \bar{\theta}^i) = \int du \mathcal{V}_0(t_A, \theta^+, \bar{\theta}^+, u^\pm) \Big|_{\theta^\pm = \theta^i u_i^\pm, \bar{\theta}^\pm = \bar{\theta}^i u_i^\pm}$$

which resolves the constraints (17), (18) for the singlet $U(1)$ part $\mathcal{X}_0 \equiv \text{Tr} (\mathcal{X})$.

The action (15) is invariant under the $\mathcal{N}=4$ superconformal group $D(2, 1; \alpha)$ with $\alpha = -\frac{1}{2}$. To show this we should use the $D(2, 1, \alpha)$ transformation laws given in [12, 20], in particular that of conformal supersymmetry,

$$\delta' \mu_H = \mu_H \left(2\Lambda - \frac{1+\alpha}{\alpha} \Lambda_0 \right), \quad \delta' \mu_A^{(-2)} = 0$$

with $\Lambda = 2i\alpha(\bar{\eta}^- \theta^+ - \eta^- \bar{\theta}^+)$, $\Lambda_0 = 2\Lambda - D^{--} \Lambda^{++}$, $\Lambda^{++} = D^{++} \Lambda$. The involved $d=1$ superfields are transformed as follows:

$$\delta' \mathcal{X} = -\Lambda_0 \mathcal{X}, \quad \delta' \mathcal{Z}^+ = \Lambda \mathcal{Z}^+, \quad \delta' V^{++} = 0.$$

The variation of the action (16) is vanishing only at $\alpha = -\frac{1}{2}$, whereas the constraints (17), (18), (21), as well as the actions (19), (20), are superconformally invariant for an arbitrary parameter α . It is important that just the field multiplier $\mathcal{V}_0$ in the action (20) provides this invariance due to its transformation law $\delta' \mathcal{V}_0 = -2\Lambda \mathcal{V}_0$ [12]. Note that at $\alpha = -1/2$ the supergroup $D(2, 1; \alpha)$ is isomorphic to $OSp(4|2)$ [22, 23], so our gauge approach in the $\mathcal{N}=4$ case implies a different $\mathcal{N}=4$ superconformal group as compared to the more customary $SU(1, 1|2)$ used e.g. in [9].

The local $U(n)$ transformations leaving the action (15) invariant are given by

$$\begin{aligned} \mathcal{X}' &= e^{i\lambda} \mathcal{X} e^{-i\lambda}, & \mathcal{Z}'^+ &= e^{i\lambda} \mathcal{Z}^+, \\ V^{++\prime} &= e^{i\lambda} V^{++} e^{-i\lambda} - i e^{i\lambda} (D^{++} e^{-i\lambda}), \end{aligned} \quad (22)$$

¹ Besides the covariant derivative $\mathcal{D}^{++}$ which commutes with $D^+, \bar{D}^+$ and so preserves the analyticity, one can define the derivative $\mathcal{D}^{--} = D^{--} + iV^{--}$, so that $[\mathcal{D}^{++}, \mathcal{D}^{--}] = D^0$ and D^0 is the operator counting the external $U(1)$ charges of superfields. The non-analytic connection V^{--} is expressed through V^{++} from this commutation relation [19].

where $\lambda_a^b(\zeta, u^\pm) \in u(n)$ is the “hermitian” analytic matrix parameter, $\hat{\lambda} = \lambda$. Using this gauge freedom we can choose the WZ gauge

$$V^{++} = -2i\theta^+\bar{\theta}^+A(t_A). \quad (23)$$

In this gauge we have

$$\begin{aligned} \mathcal{D}^{\pm\pm} &= D^{\pm\pm} + 2\theta^\pm\bar{\theta}^\pm A, \\ \mathcal{D}^- &= D^- - 2\bar{\theta}^- A, \quad \bar{\mathcal{D}}^- = \bar{D}^- - 2\theta^- A \end{aligned} \quad (24)$$

and the constraints (17), (18) are solved by

$$\mathcal{X} = X + \theta^-\bar{\theta}^-N^{++} + \theta^-\Psi^+ + \bar{\theta}^-\bar{\Psi}^+ + \dots, \quad (25)$$

where $N^{++} = N^{ik}u_i^+u_k^+$, $\Psi^+ = \Psi^iu_i^+$, $\bar{\Psi}^+ = \bar{\Psi}^iu_i^+$ and the fields $X(t_A)$, $N^{ik} = N^{(ik)}(t_A)$, $\Psi^i(t_A)$, $\bar{\Psi}^i(t_A)$ are ordinary $d=1$ fields having no dependence of the harmonics. All other fields in (25) are expressed through these fields and their covariant derivatives $\nabla_{t_A}X = \partial_{t_A}X + i[A, X]$, etc. The solution of the constraints (21) is

$$\mathcal{Z}^+ = Z^+ + \theta^+\varphi + \bar{\theta}^+\phi + 2i\theta^+\bar{\theta}^+Z^-, \quad (26)$$

where $Z^+ = Z^i(t_A)u_i^+$, $Z^- = \nabla_{t_A}Z^i(t_A)u_i^-$.

Inserting the expressions (25), (26) in the action (15) and eliminating the fields N^{ik} , ϕ , $\bar{\phi}$, φ , $\bar{\varphi}$ by their equations of motion we obtain, in the WZ gauge,

$$S_4 = S_b + S_f, \quad (27)$$

$$\begin{aligned} S_b &= \int dt \left[\text{Tr}(\nabla X \nabla X + cA) + \frac{n}{8}(\bar{Z}^{(i}Z^{k)}) (\bar{Z}_iZ_k) \right. \\ &\quad \left. + \frac{i}{2}X_0(\bar{Z}_k\nabla Z^k - \nabla\bar{Z}_kZ^k) \right], \end{aligned} \quad (28)$$

$$\begin{aligned} S_f &= -i\text{Tr} \int dt (\bar{\Psi}_k\nabla\Psi^k - \nabla\bar{\Psi}_k\Psi^k) \\ &\quad - \int dt \frac{\Psi_0^{(i}\bar{\Psi}_0^{k)}(\bar{Z}_iZ_k)}{X_0}, \end{aligned} \quad (29)$$

where

$$X_0 \equiv \text{Tr}(X), \quad \Psi_0^i \equiv \text{Tr}(\Psi^i), \quad \bar{\Psi}_0^i \equiv \text{Tr}(\bar{\Psi}^i).$$

Let us consider the bosonic limit of S_4 , i.e. the action (28). We can impose the gauge $X_a^b = 0$, $a \neq b$, using the residual invariance of WZ gauge (23): $X' = e^{i\lambda}X e^{-i\lambda}$, $Z'^k = e^{i\lambda}Z^k$, $A' = e^{i\lambda}A e^{-i\lambda} - i e^{i\lambda}(\partial_t e^{-i\lambda})$ where $\lambda_a^b(t) \in u(n)$ are ordinary $d=1$ gauge parameters. As a result of this, and after eliminating A_a^b , $a \neq b$, by the equations of motion, the action (28) takes the following form (instead of Z_a^i we introduce the new fields $Z_a^i = (X_0)^{1/2}Z_a^i$ and omit the primes on these fields),

$$\begin{aligned} S_b &= \int dt \left\{ \sum_a \dot{x}_a \dot{x}_a + \frac{i}{2} \sum_a (\bar{Z}_k^a \dot{Z}_a^k - \dot{\bar{Z}}_k^a Z_a^k) + \right. \\ &\quad \left. + \sum_{a \neq b} \frac{\text{Tr}(S_a S_b)}{4(x_a - x_b)^2} - \frac{n \text{Tr}(\hat{S}\hat{S})}{2(X_0)^2} \right\}. \end{aligned} \quad (30)$$

Here, the fields Z_a^k are subject to the constraints ²

$$\bar{Z}_i^a Z_a^i = c \quad \forall a, \quad (31)$$

and carry the residual abelian gauge $[U(1)]^n$ symmetry, $Z_a^k \rightarrow e^{i\varphi_a} Z_a^k$, with local parameters $\varphi_a(t)$. In (30) we use the following notation:

$$(S_a)_i{}^j \equiv \bar{Z}_i^a Z_a^j, \quad (32)$$

$$(\hat{S})_i{}^j \equiv \sum_a \left[(S_a)_i{}^j - \frac{1}{2} \delta_i^j (S_a)_k{}^k \right]. \quad (33)$$

Note that at $c = 0$ the constraint (31) implies $Z_a^i = 0$, i.e. a non-trivial interaction exists only for $c \neq 0$ as in the previous cases. The new feature of the $\mathcal{N}=4$ case is that not all out of the bosonic variables Z_a^i are eliminated by fixing gauges and solving the constraint; there survives a non-vanishing WZ term for them in eq. (30). After quantization these variables become purely internal (U(2)-spin) degrees of freedom.

In the Hamiltonian approach, the kinetic WZ term for Z in (30) gives rise to the following Dirac brackets:

$$[\bar{Z}_i^a, Z_b^j]_D = i\delta_b^a \delta_i^j. \quad (34)$$

With respect to these brackets the quantities (32) for each index a form $u(2)$ algebras

$$[(S_a)_i{}^j, (S_b)_k{}^l]_D = i\delta_{ab} \left\{ \delta_i^l (S_a)_k{}^j - \delta_k^j (S_a)_i{}^l \right\}. \quad (35)$$

The quantities (33) are time-independent Noether charges for the SU(2) invariance of the system (30), so the numerator of the term $\sim (X_0)^{-2}$ in (30) is a constant on the equations of motion for $Z_a^i, \bar{Z}_i^a$. So, as opposed to the $\mathcal{N}=1, 2$ cases, the $\mathcal{N}=4$ action contains a conformal potential even in the center-of-mass sector (like in [9, 11]). Modulo this extra conformal potential (last term in (30)), the bosonic limit of the $\mathcal{N}=4$ system constructed is none other than the integrable U(2)-spin Calogero model [17] in the formulation of [3, 18].

While the coordinate X_0 decouples in the bosonic limit, when all fermions are discarded, this is not the case for the full action because of the term $\sim X_0^{-1}$ in (29). The full SU(2) current contains extra fermionic terms, and its bosonic part (33) is not conserved by itself.

5. Outlook. In this paper we proposed a new gauge approach to the construction of superconformal Calogero-type systems as a superextension of the bosonic construction of [13]. The characteristic features of this approach are the presence of auxiliary supermultiplets with WZ type actions, the built-in superconformal invariance and the emergence of the Calogero coupling constant as a

² Here and in (32) we do not sum over the repeated index a .

strength of the FI term of the $U(1)$ gauge (super)field. Here we used the $U(n)$ gauging and obtained new superextensions of the A_{n-1} Calogero model and of its $U(2)$ -spin extension (in the $\mathcal{N}=4$ case). Superextensions of other conformal Calogero models could be presumably obtained by choosing other gauge groups and/or representations for the matrix and WZ superfields. Superextensions of non-conformal models can be constructed by adding proper gauge invariant (but not conformally invariant) potentials to the original superfield actions.

While in the $\mathcal{N}=1$ and $\mathcal{N}=2$ cases there is almost no freedom in the choice of the original gauged action (provided that it is required to be minimal and superconformal), it is not so in the $\mathcal{N}=4$ case due to the diversity of the $\mathcal{N}=4, d=1$ multiplets. For instance, any sort of the $\mathcal{N}=4, d=1$ multiplet has its “mirror” in which another $SU(2)$ from the full R-symmetry group $SO(4)$ of the $\mathcal{N}=4, d=1$ superalgebra is manifest. We are planning to consider these possibilities elsewhere.

In the $\mathcal{N}=4$ case we used as the basic matrix superfield the new non-abelian version of the multiplet $(\mathbf{1}, \mathbf{4}, \mathbf{3})$ defined by the constraints (17) and (18). Its simplest, quadratic action is invariant under the superconformal group $D(2, 1; \alpha)$ with $\alpha = -1/2$ (which is isomorphic to $OSp(4|2)$). It is worth noting that our gauging procedure is compatible as well with other $\mathcal{N}=4$ superconformal groups [20, 21, 22, 23]. For any value of $\alpha \neq 0$ the superconformal $\mathcal{N}=4$ gauged action has the generic form of (15) with the same S_{WZ} and S_{FI} , the only difference being in the form of the action for $\mathcal{X}$,

$$S_{\mathcal{X}}^{\alpha \neq 0} = \alpha \int \mu_H \left[\text{Tr} (\mathcal{X}^2) \right]^{-\frac{1}{2\alpha}}. \quad (36)$$

It is important that $\mathcal{X}$ is a matrix and, therefore, this action is non-trivial even in the case of $\alpha = -1$ as opposed to the standard case of the abelian $(\mathbf{1}, \mathbf{4}, \mathbf{3})$ multiplet. The second possibility at $\alpha = -1$ is to consider the matrix version of the standard conformal action

$$\tilde{S}_{\mathcal{X}}^{\alpha=-1} = \int \mu_H \text{Tr} (\mathcal{X} \ln \mathcal{X}). \quad (37)$$

It seems, however, that all such actions except for the case of $\alpha = -1/2$ yield non-trivial sigma-model type kinetic terms for the X fields, so the corresponding bosonic limits are some more general conformal models.

Acknowledgements. We thank Francois Delduc and Armen Nersessian for the interest in this work. We acknowledge a support from a grant of the Heisenberg–Landau Programme, RFBR grants 06-02-16684, 08-02-90490 and INTAS grant 05-1000008-7928 (S.F. & E.I.) and a DFG grant, project No 436 RUS/113/669 (E.I. & O.L.).

- [2] M.A. Olshanetsky, A.M. Perelomov, Phys. Rept. **71**, 313 (1981); **94**, 313 (1983).
- [3] A.P. Polychronakos, J. Phys. **A39**, 12793 (2006), [arXiv:hep-th/0607033](#).
- [4] G.W. Gibbons, P.K. Townsend Phys. Lett. **B454**, 187 (1999), [arXiv:hep-th/9812034](#).
- [5] D.Z. Freedman, P.F. Mende, Nucl. Phys. **B344**, 317 (1990).
- [6] N. Wyllard, J. Math. Phys. **41**, 2826 (2000), [arXiv:hep-th/9910160](#).
- [7] S. Bellucci, A. Galajinsky, S. Krivonos, Phys. Rev. **D68**, 064010 (2003), [arXiv:hep-th/0304087](#).
- [8] S. Bellucci, A.V. Galajinsky, E. Latini, Phys. Rev. **D71**, 044023 (2005), [arXiv:hep-th/0411232](#).
- [9] A. Galajinsky, O. Lechtenfeld, K. Polovnikov, Phys. Lett. **B643**, 221 (2006), [arXiv:hep-th/0607215](#); JHEP **0711**, 008 (2007), [arXiv:0708.1075 \[hep-th\]](#); *N=4 mechanics, WDVV equations and roots*, [arXiv:0802.4386 \[hep-th\]](#).
- [10] S. Bellucci, S. Krivonos, A. Sutulin, Nucl. Phys. **B805**, 24 (2008), [arXiv:0805.3480 \[hep-th\]](#).
- [11] S. Krivonos, O. Lechtenfeld, K. Polovnikov, *N=4 superconformal n-particle mechanics via superspace*, [arXiv:0812.5062 \[hep-th\]](#).
- [12] F. Delduc, E. Ivanov, Nucl. Phys. **B753**, 211 (2006), [arXiv:hep-th/0605211](#); **B770**, 179 (2007), [arXiv:hep-th/0611247](#); **B787**, 176 (2007), [arXiv:0706.0706 \[hep-th\]](#); Phys. Lett. **B654**, 200 (2007), [arXiv:0706.2472 \[hep-th\]](#).
- [13] A.P. Polychronakos, Phys. Lett. **B266**, 29 (1991).
- [14] A. Gorsky, N. Nekrasov, Nucl. Phys. **B414** 213 (1994), [arXiv:hep-th/9304047](#); *ibid.* **B436** 582 (1995), [arXiv:hep-th/9401017](#); Theor. Math. Phys. **100** 874 (1994).
- [15] A.P. Polychronakos, JHEP **0104**, 011 (2001), [arXiv:hep-th/0103013](#).
- [16] B. Morariu, A.P. Polychronakos, JHEP **0107**, 006 (2001), [arXiv:hep-th/0106072](#); Phys. Rev. **D72**, 125002 (2005), [arXiv:hep-th/0510034](#).
- [17] J. Gibbons, T. Hermesen, Physica **D11**, 337 (1984); S. Wojciechowski, Phys. Lett. **A111**, 101 (1985).
- [18] A.P. Polychronakos, Nucl. Phys. **B543**, 485 (1999), [arXiv:hep-th/9810211](#); Phys. Rev. Lett. **89**, 126403 (2002), [arXiv:hep-th/0112141](#).
- [19] A.S. Galperin, E.A. Ivanov, V.I. Ogievetsky, E.S. Sokatchev, *Harmonic superspace*, Cambridge University Press, 2001, 306 p.
- [20] E. Ivanov, O. Lechtenfeld, JHEP **0309**, 073 (2003), [arXiv:hep-th/0307111](#).
- [21] E. Ivanov, S. Krivonos, O. Lechtenfeld, JHEP **0303**, 014 (2003), [arXiv:hep-th/0212303](#).
- [22] L. Frappat, A. Sciarrino, P. Sorba, *Dictionary on Lie Superalgebras*, [arXiv:hep-th/9607161](#).
- [23] A. Van Proeyen, *Tools for supersymmetry*, [arXiv:hep-th/9910030](#).

[1] F. Calogero, J. Math. Phys. **10**, 2191 (1969); **10**, 2197 (1969); **12**, 419 (1971).